

Vol. 3, No. 1

**Pelvic Fractures:  
Emergency Care  
to Rehabilitation**

Postoperative Care  
of the Bariatric  
Patient



Earn CE Credits

## **Pelvic Fractures: Emergency Care to Rehabilitation**


**By Jan Foster, PhD(c), RN, CCRN**

Pelvic fractures are devastating injuries that are associated with a number of complications that often require extensive rehabilitation. Pelvic fractures represent about 0.3% to 6% of all fractures and occur in 20% of all polytrauma cases.<sup>1</sup> High-velocity trauma accounts for most pelvic fractures, including motor vehicle accidents, motor vehicle-pedestrian accidents, crush injuries, and falls.<sup>2</sup> A bimodal distribution of age is seen in pelvic trauma, as most injuries occur between the ages of 15 to 30 and 50 to 70 years. The overall average age is reportedly 31.5 years. Men account for more pelvic fractures than women, representing about 57% to 75% of all pelvic injuries.<sup>1-4</sup> Reported mortality rates range from 6.4% to 30%,<sup>5</sup> depending on the type of pelvic fracture and extrapelvic injuries and complications.

Rapid assessment and diagnosis during the emergent period, along with rehabilitation that begins as soon as the individual stabilizes, are vital to good outcomes. This article will describe types of pelvic fractures, associated intra- and extrapelvic injuries and complications, and management strategies across the continuum of emergency care and rehabilitation.

## Pelvic Anatomy

Three major bones compose the pelvis, ilium, ischium, and pubis. The ilium is situated superiorly. The uppermost portion forms the iliac crest. The right and left ilium form the pelvic girdle and articulate with the sacrum posteriorly to form the sacroiliac joint. The ischium lies inferiorly and posteriorly. The superior and inferior rami join the pubis, which lies inferiorly and anteriorly. Eventually, the three bones fuse into one at the acetabulum, which forms a socket for the head of the femur. Anterior fusion of the right and left pubi form the symphysis pubis. The circumference of the pelvic brim is shaped by the oblique plane across the sacrum on the posterior side, symphysis pubis on the anterior side, and several lateral points along the ilium. The greater pelvis, sometimes called the false pelvis, lies above this landmark. Its borders include the lumbar vertebrae, ilium laterally, and abdominal wall anteriorly. The lesser or true pelvis is below the pelvic brim, bounded by the sacrum and coccyx, inferior portions of the ilium and ischium, and pubic bones (Figure 1).<sup>6</sup>



The pelvis houses several large vascular structures. The common iliac artery branches off the abdominal aorta. The internal iliac artery branches off the common iliac and supplies most of the blood supply to the pelvic wall and viscera via several tributaries, including the superior vesicle, obturator, rectal, uterine, vaginal, pudendal, and superior and inferior gluteal arteries. The external iliac artery traverses the brim of the lesser pelvis then becomes the femoral artery as the vessel passes through the leg. The lumbar and sacral arteries, which divide off the aorta, also lie in the pelvic cavity. The veins in the pelvis, for the most part, correspond to the arteries. The internal and external iliac veins fuse to form the common iliac vein. The internal iliac vein receives blood from the pelvic organs, including the uterus, vagina, urinary bladder, rectum, prostate gland, and vas deferens as well as blood from the medial thigh and gluteal muscles. The external iliac vein continues off the femoral vein and empties blood from the anterior abdominal wall and lower extremities.<sup>6</sup>

Other vital structures within the pelvis include the reproductive organs, sigmoid colon and rectum, bladder, ureters, and urethra. Important nervous system structures that traverse the pelvis include the sacral plexus, which is composed of the 4th and 5th lumbar nerves and sacral nerves 1 through 3. Also, the femoral, sciatic, and obturator nerves pass through the pelvis.<sup>6</sup>

## Classification of Pelvic Fractures

Pelvic fractures are classified in several ways. The simplest method evaluates the nature of a pelvic-ring fracture and presence of an acetabular fracture. The pelvic-ring fracture is classified as anteroposterior compression, vertical shear, or lateral compression.<sup>7</sup> Pelvic-ring stability is a second method. Because anterior structures, such as the symphysis pubis and pubic rami, provide about 40% of pelvic rigidity, this classification system is based on posterior stability at the sacroiliac fusion. In Type A fractures, the pelvic ring is stable. Gansslen and colleagues<sup>1</sup> reported that, of 3260 pelvic injuries, 54.8% were Type A fractures. Ninety-one per cent of Type A fractures affected the anterior pelvic or iliac rim. Type B fractures are rotationally unstable. Lateral compression and complete separation at the symphysis pubis ("open-book" fractures) account for most Type B injuries and are caused by internal and external rotational forces. Concomitant acetabular fractures are commonly associated with Type B fractures.<sup>1</sup> In Type C injuries, the posterior sacroiliac articulation is completely disrupted. Seventy percent to 80% of all pelvic fractures are Type A and B (Table 1).<sup>8</sup>

Communication with adjacent tissue forms the basis for a third approach to the classification of pelvic injuries. Reportedly 2.7% of all pelvic fractures are open fractures.<sup>2</sup> Although relatively rare, they are associated with much higher morbidity and mortality than closed fractures. Open fractures most commonly result from motor-vehicle crashes. In one study, more individuals with open fractures were injured on a motorcycle (27% vs. 6%,  $p < 0.001$ ), were about 9 years younger (30 vs. 39,  $p < 0.001$ ), and were more likely to be male (75% vs. 57%,  $p < 0.02$ ), when compared to people with closed pelvic fractures.<sup>4</sup> Bilateral pubic rami fractures account for most open fractures.<sup>9</sup> Open fractures are classified as Zone I, II, or III. In Zone I fractures, communication is between the perineum, anterior pubis, medial buttock, or posterior sacrum. Zone II injuries involve the medial thigh or groin crease. Zone III injuries include communication with the posterolateral buttock or iliac crest. The most frequent type of open fractures are Zone I injuries.<sup>4</sup> Many experts recommend a temporary diverting colostomy for this type of injury to reduce the risk of fecal contamination and infectious complications.

Type	Pelvic Ring	Percentage of Injuries (%)
A	Stable	54.8
B	Partially stable, Rotationally Unstable, "Open Book", "Bucket Handle"	24.7
C	Unstable, Disruption of sacroiliac joint	20.5

## Associated Injuries and Complications

Most pelvic trauma deaths are due to associated injuries and complications rather than the pelvic fracture itself. They increase mortality from 10.8% to 31.1%.<sup>1</sup> In addition to increased mortality, associated injuries and complications account for greater blood transfusion requirements, longer hospital stays, and, predictably, increased costs of care.<sup>3</sup> Concomitant injuries can be intrapelvic or extrapelvic in nature and include intraabdominal and pelvic organ trauma, vascular injury, head injury and other neurologic injury, and other fractures. Complications during the emergent phase of trauma and hospitalization contribute to increased morbidity and mortality. Hemorrhage is the most common complication during the immediate period, whereas sepsis and multiorgan dysfunction syndrome (MODS) account for most deaths within two weeks after injury.<sup>2,10</sup>

## Abdominal and pelvic visceral injuries

Abdominal and pelvic organs often injured include the diaphragm, spleen, liver, intestine, bladder, and urethra.<sup>9,10</sup> Injuries to the ovaries, labia majora, anus, and rectum are also reported.<sup>2</sup> These injuries generally require immediate diagnosis, control of hemorrhage, and surgical or radiological repair.<sup>9,10</sup>

## Coagulopathy and vascular injury

Hemorrhage is the most significant problem associated with pelvic fracture. However, only a few individuals have osseous bleeding as a source of hemodynamic instability. Instead, hemorrhage results from intraabdominal and pelvic visceral-organ trauma and injured vasculature structures contained in the pelvis. Angiography to determine vessel injury is recommended in the presence of hypotension and hemodynamic instability when peritoneal lavage is negative for bleeding from the spleen, liver, and other intraabdominal structures. In one study, 65% of unstable patients with known pelvic fractures had pelvic-vessel injury on angiography.<sup>11</sup> In another report, angiography in 35 subjects revealed that 57% had multiple bleeding sites. Damage to the internal iliac artery and its branches was associated with unstable posterior pelvic fractures, whereas pudendal or obturator artery bleeding was associated with lateral compression fractures.<sup>12</sup> Management of hemorrhage includes aggressive fluid resuscitation, transfusions of packed red blood cells and platelets, and fresh frozen plasma and cryoprecipitate to replace clotting factors.<sup>2</sup> When hemorrhage is due to a damaged artery, embolization is sometimes used to avert bleeding, making angiography useful not only as a diagnostic tool but as a therapeutic intervention.<sup>10</sup>

## Neurologic injuries

Because pelvic fractures generally result from high-impact trauma, people often sustain open or closed head injuries during the crash. Death due to head injury usually occurs early after impact.<sup>3</sup> If the patient survives, stabilization of intracerebral hemorrhage and increased intracranial pressure, however, take priority over repair of the pelvic fracture, which may be delayed for days or weeks. Other neurologic injuries, i.e., spinal-cord,

spinal nerve-root, and sciatic-nerve injuries, occur most often with unstable pelvic-ring fractures. In a study of 83 subjects with unstable pelvic-rim fractures, 21% sustained nerve damage. Thirty-seven per cent of those had only sensory deficits, whereas 63% suffered both sensory and motor deficits. Early reduction and stabilization of unstable pelvic-ring fractures contribute to faster recovery and better prognoses. However, people with injuries at L5 are less likely to gain full function.<sup>13</sup>

## Extrapelvic fractures

It is unlikely for bones outside of the pelvis to escape injury when trauma results from motor-vehicle crashes and high-impact falls. Other fractures include bones of the upper and lower extremities, vertebrae, face, skull, sacrum, and ribs.<sup>2</sup>

## Complications

People with unstable pelvic fractures are more likely to have complications than those with stable fractures.<sup>3</sup> In a study of 236 patients with pelvic fractures, 77 (32.6%) developed 137 complications.<sup>3</sup> The most common complications included thromboembolic events, infection, and pulmonary complications.

## Thromboembolic complications

Thromboembolic events are the most common avoidable causes of morbidity and mortality in trauma patients. Deep-vein thrombosis (DVT) occurs in 35% to 60% of patients with pelvic trauma and pulmonary embolism in 2% to 10%, with a mortality of 0.5% to 4%.<sup>14</sup> The three factors leading to thrombosis and pulmonary emboli are endothelial damage, venous stasis, and hypercoagulability. All three present with pelvic fractures. There is direct injury to the pelvic veins, release of vasoactive mediators that cause venous distention, and prolonged periods of immobility and venous stasis. Both proximal thrombosis and DVT are associated with pulmonary emboli. In proximal thrombosis, thrombi most often develop in the external and iliac veins, whereas DVT involves more distal branches in the legs.

Prophylaxis of DVT is key to preventing pulmonary emboli. Standard measures include subcutaneous heparin or low-molecular-weight heparins, pneumatic compression devices on the lower extremities, and adequate hydration.<sup>14</sup> Routine screening with Doppler ultrasound and placement of vena caval filters are sometimes used to prevent pulmonary emboli. Treatment options for DVT and pulmonary emboli are somewhat limited, especially for the person with an associated intracerebral or uncontrollable hemorrhage in the chest, abdomen, pelvis, or retroperitoneum. Continuous intravenous infusion of heparin or subcutaneous injections of low-molecular-weight heparin are the treatments of choice. Vena caval filters are used in those for whom aggressive anticoagulation is contraindicated.<sup>14</sup> Thrombolytic therapy for pulmonary embolism remains controversial.<sup>15</sup>

## Infection

Various sources of infection are associated with pelvic trauma, including perineal wound infection and abscess, osteomyelitis, pneumonia, urinary tract infection, muscle and skin necrosis, fungemia, and sepsis. Infectious complications often lead to MODS, necessitating a prolonged stay in the intensive care unit. Management of focal wound infections includes repeated operative debridement, wound irrigation, split thickness skin grafts, and diverting colostomy to prevent fecal contamination. Broad-spectrum, systemic antibiotics and antifungal agents are indicated for other infectious sources. Antibiotics are adjusted according to culture results.<sup>2,4</sup>

## Pulmonary complications

Pulmonary complications are often associated with pelvic injuries. Pneumothorax should be suspected with concomitant rib fractures. Adult respiratory distress syndrome results from aspiration during the resuscitative phase of injury, pneumonia that develops later, or a septic or systemic inflammatory response.<sup>3</sup> Chest tubes, mechanical ventilation, anti-infective agents, careful management of fluids, and nutritional support are important aspects of care for pulmonary complications that accompany pelvic trauma.<sup>2</sup>

## Long-term outcomes

Achieving desired outcomes for patients with pelvic fractures requires an aggressive, multidisciplinary approach throughout the continuum of care after injury. People with open fractures experience the most complications and have the greatest rehabilitation needs. Overall, they have a poorer recovery rate than those with closed fractures. Chronic disability, impaired role performance, and poor physical function is reported.<sup>4</sup> Less often, long-term complications may include fecal and urinary incontinence, impotence, dyspareunia, and non-healing fractures.<sup>2</sup>

For people with closed fractures, early fixation is recommended, if possible. Fixation of acetabular fractures within 24 hours of injury has demonstrated a lower incidence of MODS ( $p < 0.006$ ) and decreased length of hospital stay ( $p < 0.026$ ). Patients are more likely to be discharged home rather than to rehabilitation or skilled care ( $p < 0.05$ ) and have greater functional outcomes and improved mobility.<sup>10</sup> In a study of unstable pelvic-ring fractures repaired with open reduction and internal fixation, 76% of patients returned to work, 62% full-time. Sickness Impact Profile Scores (SIP) at one year after injury indicated that 77% of these patients had only a mild disability, while 23% had a moderate disability.<sup>16</sup> These patients were allowed no weight-bearing for 3 months and required aggressive rehabilitation and assistance with activities of daily living.

## Management

Management of pelvic trauma begins with careful evaluation by emergency personnel of the circumstances surrounding the event. A number of essential facts should be determined initially, e.g., the type of incident, on which side of the vehicle the individual was sitting during a motor vehicle accident, and the speed of impact. Answers to these questions will direct appropriate diagnostic testing and rapid intervention. Full-body X-rays, blood chemistries, determination of hemoglobin, hematocrit, and platelets, coagulation studies, and urinalysis should be done to evaluate bleeding and organ injury. Other helpful diagnostic tests are peritoneal lavage and computed tomography scans.<sup>2,10</sup> Emergency management includes efforts to control hemorrhage and support hemodynamic stability. Administration of fluids and blood products should be aggressive. Angiography and possibly embolization are indicated if bleeding remains uncontrolled. Broad-spectrum antibiotics should be initiated. Orthopedic consultation and other appropriate specialists should be notified immediately. Pelvic fractures should be stabilized and fixed as early as hemodynamic and neurologic stability allows.

## Fracture stabilization

Patients who sustain pelvic fractures fall into one of two categories: those who stabilize after initial fluid resuscitation and the administration of blood products; and those who fail to respond to resuscitative efforts and remain hemodynamically unstable. Patients in the first group are managed with elective orthopedic surgery in 2 to 3 days after the injury. Stable pelvic fractures are treated with bed rest, until the patient's overall condition allows ambulation. Weight bearing may begin early in the recovery period.<sup>17</sup> Repair of unstable fractures is determined by location of the fracture, i.e., whether it is anterior or posterior. Anterior fractures can be repaired by using an external or internal approach. Injuries involving separation of the symphysis pubis or open-book fractures may be reduced and stabilized by "closing the book" with an anterior external fixation frame, which is positioned around the pelvis and held in place with pins on either side. Anterior external fixation can usually be done at the bedside under local anesthesia. Internal repair is accomplished with plates and screws, which requires general anesthesia. Unstable posterior pelvic fractures are more complex, with multiple combinations of injuries involving different bony structures in the posterior pelvis. Fracture dislocation requires open reduction and correction of bone misalignment under general anesthesia, followed by a variety of plate-and-screw fixation techniques for stabilization.<sup>17</sup>

Patients in the second category are seldom candidates for acute internal fixation. Management priorities include restoration of hemodynamic stability through angiographic embolization, laparotomy to explore and repair sources of bleeding, and external pelvic-fracture fixation. External fixation helps to control hemorrhage by reducing the volume of the pelvis, providing a tamponade effect on deep hemorrhage within the pelvis, and promote clotting.<sup>17</sup>

Open fractures are the most devastating and present the greatest challenge to the trauma team. The patient should go immediately to the operating room for exploration, control of bleeding, and repair of soft-tissue injuries.<sup>17</sup> A diverting colostomy may be done at this time to prevent stool contamination of the pelvis and subsequent sepsis.

## Nursing Care

Emergency and critical-care nurses need to be skillful in their assessment of hemodynamic monitoring, determinations of organ perfusion and oxygenation, and knowledgeable about coagulopathy, blood-product administration, and signs of hypovolemic and septic shock. Expert care of the many concomitant injuries and multisystem complications is vital to recovery in the immediate phase after injury.<sup>18</sup> In addition, several needs specific to pelvic injury warrant close attention during this time.

Anterior external fixation devices remain in place for about 6 to 8 weeks. Pins and insertion sites require frequent cleaning to prevent infection. After open fracture repair, daily wound debridement and irrigation in the operating room is recommended for at least 4 days. Frequent dressing changes in the intensive care unit are important to reduce the risk of sepsis.<sup>17</sup> The dressing may be held in place with an abdominal binder, as there will generally be an open, gaping wound, making it otherwise difficult to secure the dressings. The abdominal binder may also facilitate coughing and deep breathing and promote greater patient movement, which will help to prevent pneumonia. Tissue integrity of the colostomy should be assessed frequently due to the risk of compromised blood flow during the initial injury. A colostomy bag should be secured over the stoma with skin-protective sealant. Urinary output, an important indicator of intravascular volume and organ perfusion, should be monitored closely. Hematuria should be noted, as it may signify renal or bladder injury. The urinary catheter should be stabilized with a catheter-securing device to prevent further trauma to the urinary meatus and bladder.

Pain management is a high priority during this time, as orthopedic injury is one of the most severe types of pain. Continuous intravenous infusion of pain medication provides a steady blood level and is generally more effective in maintaining analgesia at lower overall doses.<sup>18,19</sup>

The patient is at risk for complications due to immobility during this relative time of prolonged bed rest, which is potentiated by poor tissue perfusion during the emergent phase of injury. The use of therapeutic beds, early provision of parenteral or enteral nutrition, and pneumatic-compression devices help to prevent skin breakdown and deep-vein thrombosis, respectively. Intravenous catheters and enteral feeding tubes should be secured. A bendable armboard with soft material covering can assist in securing the catheter while minimizing skin damage. Frequent site assessment and dressing changes are necessary to prevent infection. Bowel elimination is achieved with stool softeners, particularly when the patient receives high doses of narcotic for pain management.<sup>19</sup> Once the patient is well stabilized, wound healing, oral nutrition, physical therapy, and general conditioning become the main concerns.

## Conclusions

Most people with pelvic fractures survive their injury. If not, death is usually due to concomitant injuries. Control of hemorrhage, infection, diagnosis and management of

related injuries, early fixation of fractures, and early initiation of rehabilitation are critical to better functional outcomes for patients with pelvic fractures. Patients suffering pelvic injury require a coordinated effort by numerous healthcare professionals throughout the trajectory of care, beginning with the trauma team in the field, continuing with the rehabilitation team to discharge, and beyond, as outpatient rehabilitation continues for weeks or months. With early intervention and prevention of problems through collaborative efforts, patients with pelvic fractures can achieve optimal functional outcomes and return to the community.

## References

1. Gansslen A, Pohleman T, Paul C, Lobenhoffer P, Tscherne H. Epidemiology of pelvic ring fractures. *Injury* 1996;27(Suppl 1:SA):13-20.
2. Ferrera PC, Hill DA. Good outcomes of open pelvic fractures. *Injury* 1999;30(3): 187-90.
3. Poole GV, Ward EF, Muakkassa EF, Hsu HS, Griswold JA, Rhodes RS. Pelvic fracture from blunt trauma. Outcomes is determined by associated injuries. *Ann Surg* 1991;213(6):532-8.
4. Brenneman FD, Katyal D, Boulanger BR, Tile M, Redelmeier DA. Long-term outcomes in open pelvic fractures. *J Trauma* 1997;42(5):773-7.
5. Wubben RC. Mortality rate of pelvic fractures. *Wis Med J* 1996;95(10):702-4.
6. Guyton A. *Anatomy and Physiology*. Philadelphia:W. B. Saunders, 1996.
7. Plaisier BR, Meldon SW, Super DM, Malangoni MA. Improved outcomes after early fixation of acetabular fractures. *Injury* 2000;31(2):81-4.
8. Tile M. Acute pelvic fractures: causation and classification. *J Am Acad Orthop Surg* 1996;4(3):143-151.
9. Killeen KL, DeMeo JH. CT detection of serious internal and skeletal injuries in patients with pelvic fractures. *Acad Radiol* 1999;6(4):224-8.
10. Pajenda GS, Seitz H, Mousavi M, Vecsei V. Concomitant intra-abdominal injuries in pelvic trauma. *Wien Klin Wochenscher* 1998;110(23):834-40.
11. Agolini SF, Shah K, Jaffe J, Newcomb J, Rhodes M, Reed JF. Arterial embolization is a rapid and effective technique for controlling pelvic fracture hemorrhage. *J Trauma* 1997;43(3):395-400.
12. O'Neill PA, Riina J, Sclafani S, Tornetta P. Angiographic findings in pelvic fractures. *Clin Orthop* 1996;329:60-7.

13. Reilly MC, Zinar DM, Matta JM. Neurologic injuries in pelvic-ring fractures. Clin Orthop 1996;329:28-36.
14. Montgomery KD, Geerts WH, Potter HG, Helfet DL. Thromboembolic complications in patients with pelvic trauma. Clin Orthop 1996;329:68-85.
15. Scroggins N. Hemorrhagic disorders associated with thrombolytic therapy. Critical Care Nursing Clinics of North America 2000;12(3):353-363.
16. Gruen GS, Leit ME, Gruen RJ, Garrison HG, Auble TE, Peitzman AB. Functional outcomes of patients with unstable pelvic-ring fractures with open reduction and internal fixation. J Trauma 1995;39(5):838-44.
17. Cryer HG, Johnson E. Pelvic fractures. In: Feliciano D, Moore, E, Mattox K. Trauma. Stanford, CT:Appleton & Lange, 1996, 635-59.
18. Hudak CM, Gallo BM, Morton PG. Critical Care Nursing: A Holistic Approach. Philadelphia: Lippincott, 1998.
19. Boehnlein MJ, Marek JF. Postoperative nursing. In: Phipps WJ, Sands J, Marek JF. Medical Surgical Nursing Concepts and Clinical Practice. St. Louis: Mosby, 1999, 525-550.



Janet G. Whetstone Foster, RN, MSN, CCRN, is Assistant Professor of Adult Acute and Critical Care Nursing and a clinical nurse specialist in adult health at Houston Baptist University, Houston, Texas. She is President of Nursing Inquiry & Intervention Inc., a healthcare consulting firm. A sought-after speaker, she specializes in critical-care nursing, particularly the area of neuromuscular blockade.